

ON THE INDEX OF FAREY SEQUENCES

FLORIN P. BOCA, RADU N. GOLOGAN AND ALEXANDRU ZAHARESCU

ABSTRACT. We prove some asymptotic formulae concerning the distribution of the index of Farey fractions of order Q as $Q \rightarrow \infty$.

1. INTRODUCTION

Let $\mathcal{F}_Q = \{\gamma_1, \dots, \gamma_{N(Q)}\}$ denote the Farey sequence of order Q with $\gamma_1 = 1/Q < \gamma_2 < \dots < \gamma_{N(Q)} = 1$. This sequence is extended by $\gamma_{i+N(Q)} = \gamma_i + 1$, $1 \leq i \leq N(Q)$. It is well-known that

$$N(Q) = \sum_{j=1}^Q \varphi(j) = \frac{3Q^2}{\pi^2} + O(Q \log Q).$$

For any two consecutive Farey fractions $\gamma_i = a_i/q_i < \gamma_{i+1} = a_{i+1}/q_{i+1}$, one has $a_{i+1}q_i - a_iq_{i+1} = 1$ and $q_i + q_{i+1} > Q$. Conversely, if q and q' are two coprime integers in $\{1, \dots, Q\}$ with $q + q' > Q$, then there are unique $a \in \{1, \dots, q\}$ and $a' \in \{1, \dots, q'\}$ for which $a'q - aq' = 1$, and $a/q < a'/q'$ are consecutive Farey fractions of order Q . Therefore, the pairs of coprime integers (q, q') with $q + q' > Q$ are in one-to-one correspondence with the pairs of consecutive Farey fractions of order Q . Moreover, the denominator q_{i+2} of γ_{i+2} can be easily expressed (cf. [5]) by means of the denominators of γ_i and γ_{i+1} as

$$q_{i+2} = \left\lceil \frac{Q + q_i}{q_{i+1}} \right\rceil q_{i+1} - q_i.$$

This formula was recently used in the study of various statistical properties of the Farey fractions [1], [2], [4], leading to the definition in [2] of a new and interesting area-preserving transformation T of the Farey triangle

$$\mathcal{T} = \{(x, y) \in [0, 1]^2; x + y > 1\},$$

defined by

$$T(x, y) = \left(y, \left\lceil \frac{1+x}{y} \right\rceil y - x \right) = \left(y, 1 - x - yG\left(\frac{y}{1+x}\right) \right),$$

where $G(t) = \{1/t\}$ is the Gauss map on the unit interval. The set $\mathcal{T}$ decomposes as the union of disjoint sets (see Figure 1)

$$\mathcal{T}_k = \left\{ (x, y) \in \mathcal{T}; \left\lceil \frac{1+x}{y} \right\rceil = k \right\} \quad k \in \mathbb{N}^* = \{1, 2, 3, \dots\},$$

Research partially supported by ANSTI grant C6189/2000.

and

$$T(x, y) = (y, ky - x), \quad (x, y) \in \mathcal{T}_k.$$




FIGURE 1. The decomposition $\mathcal{T} = \cup_{k=1}^{\infty} \mathcal{T}_k$.

For later use, we also define the upper triangles $\mathcal{T}'_k$ with vertices at $(1, 2/k)$, $((k-1)/(k+1), 2/(k+1))$ and $(1, 2/(k+1))$, $k \geq 2$, and the lower triangles $\mathcal{T}''_k$ with vertices at $(k/(k+2), 2/(k+2))$, $((k-1)/(k+1), 2/(k+1))$ and $(1, 2/(k+1))$, $k \geq 1$.

For all integer $i \geq 0$, we have

$$T^i(x, y) = (L_i(x, y), L_{i+1}(x, y)),$$

with

$$\begin{aligned} L_{i+1}(x, y) &= \kappa_i(x, y)L_i(x, y) - L_{i-1}(x, y), \\ L_0(x, y) &= x, \quad L_1(x, y) = y \\ \kappa_i(x, y) &= \kappa_{i-1} \circ T(x, y), \quad \kappa_1(x, y) = [(1+x)/y]. \end{aligned}$$

It was noticed and extensively used in [2] that

$$T\left(\frac{q}{Q}, \frac{q'}{Q}\right) = \left(\frac{q'}{Q}, \frac{q''}{Q}\right),$$

whenever q , q' and q'' are denominators of three consecutive Farey fractions from $\mathcal{F}_Q$. This shows immediately that

$$(1.1) \quad \kappa_1\left(\frac{q_{i-1}}{Q}, \frac{q_i}{Q}\right) = \left[\frac{Q + q_{i-1}}{q_i}\right]$$

and

$$(1.2) \quad \begin{aligned} \kappa_{r+1}\left(\frac{q_{i-1}}{Q}, \frac{q_i}{Q}\right) &= \kappa_1 \circ T^r\left(\frac{q_{i-1}}{Q}, \frac{q_i}{Q}\right) = \kappa_1\left(\frac{q_{i+r-1}}{Q}, \frac{q_{i+r}}{Q}\right) \\ &= \left[\frac{Q + q_{i+r-1}}{q_{i+r}}\right], \quad r \in \mathbb{N}. \end{aligned}$$

We also note that

$$(1.3) \quad \mathcal{T}_k = \{(x, y) \in \mathcal{T} ; \kappa_1(x, y) = k\}.$$

In [4], the integer

$$(1.4) \quad \nu_Q(\gamma_i) = \left[\frac{Q + q_{i-1}}{q_i}\right] = \kappa_1\left(\frac{q_{i-1}}{Q}, \frac{q_i}{Q}\right) = \frac{q_{i+1} + q_{i-1}}{q_i} = \frac{a_{i+1} + a_{i-1}}{a_i}$$

was called the index of the fraction γ_i in $\mathcal{F}_Q$, and various new and interesting results concerning their distribution were proved, including the striking closed form formulae

$$(1.5) \quad \sum_i \nu_Q(\gamma_i) = 3N(Q) - 1$$

and

$$\begin{aligned} \sum_{q=1}^Q \# \left\{ \gamma_i = \frac{a_i}{q_i} ; q_i = q \text{ and } \nu_Q(\gamma_i) = \left[\frac{2Q+1}{q_i}\right] - 1 \right\} \\ = Q(2Q+1) - N(2Q) - 2N(Q) + 1. \end{aligned}$$

The following asymptotic formulae were also proved in [4]:

$$\sum_i \nu_Q(\gamma_i)^2 = \frac{24}{\pi^2} Q^2 \left(\log 2Q - \frac{\zeta'(2)}{\zeta(2)} - \frac{17}{8} + 2\gamma \right) + O(Q \log^2 Q),$$

$$(1.6) \quad \begin{aligned} L(Q, k) &= l_k N(Q) + O\left(k + \frac{Q \log Q}{k}\right), \\ U(Q, k) &= u_k N(Q) + O\left(k + \frac{Q \log Q}{k}\right), \end{aligned}$$

where

$$\begin{aligned} L(Q, k) &= \# \left\{ \gamma_i ; \nu_Q(\gamma_i) = k = \left[\frac{2Q+1}{q_i}\right] - 1 \right\}, \\ U(Q, k) &= \# \left\{ \gamma_i ; \nu_Q(\gamma_i) = k = \left[\frac{2Q+1}{q_i}\right] \right\}, \end{aligned}$$

and

$$\begin{aligned} l_k &= 4 \left(\frac{1}{(k+1)^2} - \frac{1}{k+1} + \frac{1}{k+2} \right), \\ u_k &= \begin{cases} 0 & \text{if } k = 1, \\ 4 \left(\frac{1}{k} - \frac{1}{k+1} - \frac{1}{(k+1)^2} \right) & \text{if } k \geq 2. \end{cases} \end{aligned}$$

In an earlier version of [4], it was proved that

$$(1.7) \quad \sum_{\gamma_i \leq t} \nu_Q(\gamma_i) = 3N(Q)t + O(Q^{3/2+\varepsilon}) \quad \text{whenever } t \in [0, 1].$$

It was also conjectured that for every $h \in \mathbb{N}^*$, there is a constant $A(h)$ such that

$$S_h(Q) = \sum_i \nu_Q(\gamma_i) \nu_Q(\gamma_{i+h}) \sim A(h)N(Q)$$

as $Q \rightarrow \infty$. In this note, we first prove that this conjecture holds, finding also the finite constant

$$A(h) = 2 \iint_{\mathcal{T}} \kappa_1(s, t) \kappa_{h+1}(s, t) ds dt = 2 \sum_{m, n=1}^{\infty} mn \operatorname{area}(T^h \mathcal{T}_m \cap \mathcal{T}_n).$$

All the numbers $A(h)$ are rational and $A(1) = \frac{192}{35} \approx 5.4857$, $A(2) = \frac{796727}{90090} \approx 8.8437$. One can also show that $A(h) \ll 1 + \log h$. It would be interesting to investigate whether $A(h)$ is bounded by an absolute constant.

We also consider the more general situation where the Farey fractions belong to a subinterval of $[0, 1]$. More precisely, we set

$$S_{h,t}(Q) = \sum_{\gamma_i \in \mathcal{F}_Q \cap [0, t]} \nu_Q(\gamma_i) \nu_Q(\gamma_{i+h}),$$

and prove

Theorem 1.1. (i) *For every integer $h \geq 1$, we have*

$$S_h(Q) = A(h)N(Q) + O_h(Q \log^2 Q)$$

as $Q \rightarrow \infty$.

(ii) *For every integer $h \geq 1$ and every $t \in [0, 1]$, we have for all $\varepsilon > 0$,*

$$S_{h,t}(Q) = tA(h)N(Q) + O_{h,\varepsilon}(Q^{3/2+\varepsilon}).$$

For $0 < \alpha < 2$, we define

$$B_\alpha = \iint_{\mathcal{T}} \left[\frac{1+s}{t} \right]^\alpha ds dt = \sum_{k=1}^{\infty} k^\alpha \operatorname{area}(\mathcal{T}_k) \ll \sum_{k=1}^{\infty} k^{\alpha-3} < \infty.$$

Employing the results from [2], we give the following generalization of (1.7):

Theorem 1.2. (i) *For every $\alpha \in (0, 2)$, we have*

$$\left| \sum_i \nu_Q(\gamma_i)^\alpha - 2N(Q)B_\alpha \right| \ll_\alpha E_\alpha(Q) = \begin{cases} Q \log Q & \text{if } \alpha < 1, \\ Q \log^2 Q & \text{if } \alpha = 1, \\ Q^\alpha \log Q & \text{if } 1 < \alpha < 2. \end{cases}$$

(ii) *For every $\alpha \in (0, 3/2)$ and $t \in [0, 1]$, we have for all $\varepsilon > 0$,*

$$\left| \sum_{\gamma_i \leq t} \nu(\gamma_i)^\alpha - 2tN(Q)B_\alpha \right| \ll_{\alpha,\varepsilon} F_\alpha(Q) = \begin{cases} Q^{3/2+\varepsilon} & \text{if } \alpha \leq 1, \\ Q^{\alpha+1/2+\varepsilon} & \text{if } \alpha > 1. \end{cases}$$

Note that

$$B_1 = \sum_{k=1}^{\infty} k \operatorname{area}(\mathcal{T}_k) = \frac{1}{6} + \sum_{k=2}^{\infty} \frac{4k}{k(k+1)(k+2)} = \frac{3}{2},$$

which is consistent with (1.5) and (1.4).

Finally, we show that (1.6) with error $O(Q \log Q/k)$ can be derived from Lemma 2 in [2]. In our framework the geometrical significance of the constants l_k and u_k is apparent, as

$$l_k = 2 \operatorname{area}(\mathcal{T}_k'') \quad \text{and} \quad u_k = 2 \operatorname{area}(\mathcal{T}_k').$$

Furthermore, if we set for $t \in (0, 1]$,

$$L(Q, k, t) = \# \left\{ \gamma_i = \frac{a_i}{q_i} \leq t; \nu_Q(\gamma_i) = k = \left\lfloor \frac{2Q+1}{q_i} \right\rfloor - 1 \right\}$$

and

$$U(Q, k, t) = \# \left\{ \gamma_i = \frac{a_i}{q_i} \leq t; \nu_Q(\gamma_i) = k = \left\lfloor \frac{2Q+1}{q_i} \right\rfloor \right\},$$

then we deduce as a result of Lemma 10 in [2] the following generalization of (1.6):

Theorem 1.3. *For every $t \in (0, 1]$ and every $\varepsilon > 0$, we have*

$$L(Q, k, t) = tl_k N(Q) + O_\varepsilon \left(\frac{Q^{3/2+\varepsilon}}{k} \right)$$

and

$$U(Q, k, t) = tu_k N(Q) + O_\varepsilon \left(\frac{Q^{3/2+\varepsilon}}{k} \right).$$

Acknowledgements We are grateful to Richard Hall and Peter Shiu for correspondence on [4] and to the anonymous referee for suggestions leading to the improvement of this paper.

2. PROOF OF THE MAIN RESULTS

We denote throughout

$$\mathbb{Z}_{\text{vis}}^2 = \{(a, b) \in \mathbb{Z}^2; \gcd(a, b) = 1\}.$$

By (1.2), we get

$$\nu_Q(\gamma_{i+h}) = \left\lfloor \frac{Q + q_{i+h-1}}{q_{i+h}} \right\rfloor = \kappa_{h+1} \left(\frac{q_{i-1}}{Q}, \frac{q_i}{Q} \right),$$

leading to

$$S_h(Q) = \sum_{\gamma_i \in \mathcal{F}_Q} \kappa_1 \left(\frac{q_{i-1}}{Q}, \frac{q_i}{Q} \right) \kappa_{h+1} \left(\frac{q_{i-1}}{Q}, \frac{q_i}{Q} \right).$$

Since the pairs of denominators of consecutive elements in $\mathcal{F}_Q$ coincide with the elements of the set

$$Q\mathcal{T} \cap \mathbb{Z}_{\text{vis}}^2 = \{(a, b) \in \mathbb{Z}_{\text{vis}}^2; a + b > Q \geq a, b \geq 1\},$$

we may write

$$S_h(Q) = \sum_{k=1}^{\infty} \sum_{(a,b) \in Q\mathcal{T} \cap \mathbb{Z}_{\text{vis}}^2} \kappa_1\left(\frac{a}{Q}, \frac{b}{Q}\right) \kappa_{h+1}\left(\frac{a}{Q}, \frac{b}{Q}\right).$$

Taking also stock on (1.3), this further yields

$$\begin{aligned} S_h(Q) &= \sum_{k=1}^{\infty} \sum_{(a,b) \in Q\mathcal{T}_k \cap \mathbb{Z}_{\text{vis}}^2} \kappa_{h+1}\left(\frac{a}{Q}, \frac{b}{Q}\right) \\ (2.1) \quad &= \sum_{k=1}^{\infty} k \sum_{l=1}^{\infty} l \# \left\{ (a, b) \in Q\mathcal{T}_k \cap \mathbb{Z}_{\text{vis}}^2; \kappa_1 T^h\left(\frac{a}{Q}, \frac{b}{Q}\right) = l \right\} \\ &= \sum_{k,l=1}^{\infty} kl \# \left\{ (a, b) \in Q\mathcal{T}_k \cap \mathbb{Z}_{\text{vis}}^2; T^h\left(\frac{a}{Q}, \frac{b}{Q}\right) \in \mathcal{T}_l \right\} \\ &= \sum_{k,l=1}^{\infty} kl \# (Q(\mathcal{T}_k \cap T^{-h}\mathcal{T}_l) \cap \mathbb{Z}_{\text{vis}}^2). \end{aligned}$$

If we set

$$\mathcal{T}_k^* = \bigcup_{n=k}^{\infty} \mathcal{T}_n,$$

then

$$(2.2) \quad \mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^* = \left(\bigcup_{m=k}^{\infty} \mathcal{T}_m \right) \cap T^{-h} \left(\bigcup_{n=l}^{\infty} \mathcal{T}_n \right) = \bigcup_{m=k}^{\infty} \bigcup_{n=l}^{\infty} (\mathcal{T}_m \cap T^{-h}\mathcal{T}_n).$$

We wish to estimate

$$A_{k,l}(Q) = \#(Q(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*) \cap \mathbb{Z}_{\text{vis}}^2).$$

Since the sets from the right-hand side of (2.2) are mutually disjoint, we now have

$$\begin{aligned} \sum_{k,l=1}^{\infty} A_{k,l}(Q) &= \sum_{k,l=1}^{\infty} \sum_{m=k}^{\infty} \sum_{n=l}^{\infty} \#(Q(\mathcal{T}_m \cap T^{-h}\mathcal{T}_n) \cap \mathbb{Z}_{\text{vis}}^2) \\ &= \sum_{m,n=1}^{\infty} \#(Q(\mathcal{T}_m \cap T^{-h}\mathcal{T}_n) \cap \mathbb{Z}_{\text{vis}}^2) \sum_{k=1}^m \sum_{l=1}^n 1 \\ &= \sum_{m,n=1}^{\infty} mn \#(Q(\mathcal{T}_m \cap T^{-h}\mathcal{T}_n) \cap \mathbb{Z}_{\text{vis}}^2), \end{aligned}$$

so that, by (2.1),

$$(2.3) \quad S_h(Q) = \sum_{k,l=1}^{\infty} A_{k,l}(Q).$$

The following lemma is proved in a similar way as Lemma 2 in [2].

Lemma 2.1. *Let $\Omega \subset [0, R_1] \times [0, R_2]$ be a region in $\mathbb{R}^2$ with rectifiable boundary $\partial\Omega$ and let $g : \Omega \rightarrow \mathbb{R}$ be a C^1 function on Ω . Suppose that $R \geq \min(R_1, R_2)$. Then we have*

$$\begin{aligned} \sum_{(a,b) \in \Omega \cap \mathbb{Z}_{\text{vis}}^2} g(a,b) &= \frac{6}{\pi^2} \iint_{\Omega} g(x,y) dx dy + O(\|Dg\|_{\infty} \text{area}(\Omega) \log R) \\ &\quad + O\left(\|g\|_{\infty} \left(R + \frac{\text{area}(\Omega)}{R} + \text{length}(\partial\Omega) \log R\right)\right), \end{aligned}$$

where

$$\|Dg\|_{\infty} = \sup_{(x,y) \in \Omega} \left| \frac{\partial g}{\partial x}(x,y) \right| + \left| \frac{\partial g}{\partial y}(x,y) \right|.$$

Corollary 2.2. *Let $\Omega \subseteq [0, R_1] \times [0, R_2]$ be a bounded region with rectifiable boundary $\partial\Omega$ and let $R \geq \min(R_1, R_2)$. Then we have*

$$\#(\Omega \cap \mathbb{Z}_{\text{vis}}^2) = \frac{6 \text{area}(\Omega)}{\pi^2} + O\left(R + \text{length}(\partial\Omega) \log R + \frac{\text{area}(\Omega)}{R}\right).$$

Remark 2.3. Let $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$, let Φ be the linear transformation on $\mathbb{R}^2$ defined by $\Phi(x,y) = (ax + by, cx + dy)$, and let Ω be a bounded region in $\mathbb{R}^2$. Then $\#(\Omega \cap \mathbb{Z}_{\text{vis}}^2) = \#(\Phi\Omega \cap \mathbb{Z}_{\text{vis}}^2)$, which implies in turn that

$$\#(Q\Omega \cap \mathbb{Z}_{\text{vis}}^2) = \#(Q(T\Omega) \cap \mathbb{Z}_{\text{vis}}^2) = \dots = \#(Q(T^h\Omega) \cap \mathbb{Z}_{\text{vis}}^2)$$

whenever $\Omega \subseteq \mathcal{T}$. In particular, this gives

$$(2.4) \quad \#(Q(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*) \cap \mathbb{Z}_{\text{vis}}^2) = \#(Q(T^h\mathcal{T}_k^* \cap \mathcal{T}_l^*) \cap \mathbb{Z}_{\text{vis}}^2).$$

Lemma 2.4. [2, Lemma 5] *Let $r \geq 1$ and suppose that $i \neq j$ and that $\max(L_i(x,y), L_j(x,y)) \leq 2^{-r-1}$ for some point $(x,y) \in \mathcal{T}$. Then $|j - i| > r + 1$.*

Corollary 2.5. *Assume that $\min(k, l) > c_h = 2^{h+1}$. Then we have*

$$\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^* = \emptyset.$$

Proof. Suppose that there exists $(x,y) \in \mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*$. Then $L_1(x,y) = y \leq 2/k < 2^{-h}$ and $L_1(T^h(x,y)) = L_{h+1}(x,y) \leq 2/l < 2^{-h}$. We now infer from Lemma 2.4 that $|h + 1 - 1| > h$, which is a contradiction. $\square$

Remark 2.6. A careful look at the proof of Lemma 2.4 shows that the constant 2^{r+1} can be lowered to $4r+2$ (see Lemma 3.4 and Remark 3.5 in [3]). Note also that $T^{\pm 1}\mathcal{T}_m^* \subset \mathcal{T}_1$ for all $m \geq 5$. By Lemma 3.4 in [3] it also follows that for all $r \geq 2$ and all $m \geq c_r = 4r+2$, we have $\cup_{i=2}^r T^{\pm i}\mathcal{T}_m^* \subset \mathcal{T}_2$. In summary, we have

$$T^h\mathcal{T}_m^* \cap \mathcal{T}_n^* = \emptyset$$

whenever $m \geq c_h$ and $n \geq 3$.

Since $Q\mathcal{T}_k^* \cap \mathbb{Z}_{\text{vis}}^2 = \emptyset$ whenever $2Q/k < 1$, Remark 2.3 shows that $A_{k,l}(Q) = 0$ unless $\max(k, l) \leq 2Q$. By Corollary 2.5, $A_{k,l}(Q) = 0$ unless $\min(k, l) \leq c_h$. Thus (2.3) provides

$$(2.5) \quad S_h(Q) = \sum_{k,l=1}^{2Q} A_{k,l}(Q) = C_h(Q) + D_h(Q),$$

where

$$C_h(Q) = \sum_{k=c_h+1}^{2Q} \sum_{l=1}^{c_h} A_{k,l}(Q) \quad \text{and} \quad D_h(Q) = \sum_{k=1}^{c_h} \sum_{l=1}^{2Q} A_{k,l}(Q).$$

Suppose first that $k > c_h$. Applying Corollary 2.2 to $\Omega = Q(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*)$ with $\text{area}(\Omega) \ll Q^2/k^2$, and with

$$\begin{aligned} \text{length}(\partial\Omega) &= Q \text{length} \left(\bigcup_{r=l}^{\infty} T^{-h}\mathcal{T}_r \cap \mathcal{T}_k^* \right) = Q \text{length} \left(\bigcup_{r=l}^{c_h} T^{-h}\mathcal{T}_r \cap \mathcal{T}_k^* \right) \\ &\leq Q \sum_{r=l}^{c_h} \text{length}(T^{-h}\mathcal{T}_r \cap \mathcal{T}_k^*) \ll_h \frac{Q}{k}, \end{aligned}$$

we infer that

$$(2.6) \quad A_{k,l}(Q) = \frac{6Q^2}{\pi^2} \text{area}(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*) + O_h \left(\frac{Q \log Q}{k} \right).$$

By (2.6) and $\text{area}(\mathcal{T}_k^*) = O(k^{-2})$ we gather

$$\begin{aligned} (2.7) \quad C_h(Q) &= \frac{6Q^2}{\pi^2} \sum_{k=c_h+1}^{2Q} \sum_{l=1}^{c_h} \text{area}(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*) + O_h(Q \log^2 Q) \\ &= \frac{6Q^2}{\pi^2} \sum_{k=c_h+1}^{\infty} \sum_{l=1}^{\infty} \text{area}(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*) + O_h(Q \log^2 Q). \end{aligned}$$

When $k \leq c_h$, we employ Remark 2.3 and equality (2.4) to get

$$A_{k,l}(Q) = \#(Q(\mathcal{T}_l^* \cap T^h\mathcal{T}_k^*)).$$

Applying now Corollary 2.2 to $\Omega = Q(\mathcal{T}_l^* \cap T^h \mathcal{T}_k^*)$ with $\text{length}(\partial\Omega) \ll_h Q/l$, and using the fact that T is area-preserving, we obtain

$$(2.8) \quad \begin{aligned} A_{k,l}(Q) &= \frac{6}{\pi^2} \text{area}(\mathcal{T}_l^* \cap T^h \mathcal{T}_k^*) + O_h\left(\frac{Q \log Q}{l}\right) \\ &= \frac{6}{\pi^2} \text{area}(\mathcal{T}_k^* \cap T^{-h} \mathcal{T}_l^*) + O_h\left(\frac{Q \log Q}{l}\right). \end{aligned}$$

We may now employ (2.7) and $\text{area}(\mathcal{T}_l^*) = O(l^{-2})$, to get

$$(2.9) \quad D_h(Q) = \frac{6}{\pi^2} \sum_{k=1}^{c_h} \sum_{l=1}^{\infty} \text{area}(\mathcal{T}_k^* \cap T^{-h} \mathcal{T}_l^*) + O_h(Q \log^2 Q).$$

Inserting (2.9) and (2.7) into (2.5), we collect

$$(2.10) \quad S_h(Q) = \frac{6}{\pi^2} \sum_{k,l=1}^{\infty} \text{area}(\mathcal{T}_k^* \cap T^{-h} \mathcal{T}_l^*) + O_h(Q \log^2 Q).$$

We conclude the proof of Theorem 1.1 (i) by noting as in (2.3) that the sum in the right-hand side of (2.10) is equal to

$$(2.11) \quad \sum_{m,n=1}^{\infty} \text{area}(T^h \mathcal{T}_m^* \cap \mathcal{T}_n^*) = \sum_{m,n=1}^{\infty} mn \text{area}(T^h \mathcal{T}_m \cap \mathcal{T}_n) = \frac{A(h)}{2}.$$

We set

$$f = f_0 + R_0 \quad \text{and} \quad g = f \circ T^{-h} = f_0 \circ T^{-h} + R_0 \circ T^{-h},$$

with

$$f_0 = \sum_{m=1}^{c_h-1} e_{\mathcal{T}_m^*}, \quad R_0 = \sum_{m=c_h}^{\infty} e_{\mathcal{T}_m^*},$$

where e_S denotes the characteristic function of the set S . By Remark 2.6 we gather that the product of R_0 and g equals

$$\sum_{m=c_h}^{\infty} \sum_{n=1}^{\infty} e_{T^h \mathcal{T}_m^* \cap \mathcal{T}_n^*} = \sum_{m=c_h}^{\infty} \sum_{n=1}^2 e_{T^h \mathcal{T}_m^* \cap \mathcal{T}_n^*}.$$

Hence,

$$\iint_T R_0(s, t) g(s, t) ds dt \leq 2 \sum_{m=c_h}^{\infty} \text{area}(\mathcal{T}_m^*) \ll \sum_{m=1}^{\infty} \frac{1}{m^2} \ll 1.$$

In a similar way, we get

$$\iint_T f(s, t) (R_0 \circ T^{-h})(s, t) ds dt \ll 1.$$

Using (2.11) we may now write

$$(2.12) \quad \begin{aligned} A(h) &= 2 \iint_T f(s, t)(f \circ T^{-h})(s, t) ds dt \\ &\ll 1 + 2 \iint_T f_0(s, t)g_0(s, t) ds dt. \end{aligned}$$

The function f_0 belongs to $L^2(\mathcal{T})$ as a result of

$$\begin{aligned} f_0 &= \sum_{m=1}^{c_h-1} \sum_{k=m}^{\infty} e_{\mathcal{T}_k} = \sum_{k=1}^{\infty} \sum_{m=1}^{\min(k, c_h-1)} e_{\mathcal{T}_k} = \sum_{k=1}^{\infty} \min(k, c_h-1) e_{\mathcal{T}_k} \\ &= \sum_{k=1}^{c_h-1} k e_{\mathcal{T}_k} + (c_h-1) \sum_{k=c_h}^{\infty} e_{\mathcal{T}_k} = \sum_{k=1}^{c_h-1} k e_{\mathcal{T}_k} + (c_h-1) e_{\mathcal{T}_{c_h}^*}, \end{aligned}$$

which gives in turn

$$\|f_0\|_{L^2(\mathcal{T})} = \sum_{k=1}^{c_h-1} k^2 \text{area}(\mathcal{T}_k) + (c_h-1)^2 \text{area}(\mathcal{T}_{c_h}^*) \ll \sum_{k=1}^{c_h} \frac{1}{k} + 1 \ll 1 + \log h.$$

Since T is area-preserving, we also have

$$\|g_0\|_{L^2(\mathcal{T})} = \|f_0 \circ T^{-h}\|_{L^2(\mathcal{T})} = \|f_0\|_{L^2(\mathcal{T})},$$

and the Cauchy-Schwarz inequality gives in conjunction with (2.12)

$$A(h) \ll 1 + 2\|f_0\|_{L^2(\mathcal{T})}\|g_0\|_{L^2(\mathcal{T})} \ll 1 + \log h.$$

To prove (ii), we proceed as in Section 8 in [2] and note that the relation $a'q - aq' = 1$ between two consecutive Farey fractions of order Q shows that $a = q - \bar{q}'$, where $\bar{q}' \in \{1, \dots, q\}$ denotes the multiplicative inverse of $q' \pmod{q}$. Thus the condition $a/q \in I$ with $I = (\alpha, \beta] \subseteq (0, 1]$ interval, is equivalent to $\bar{q}' \in I_q = [(1 - \beta)q, (1 - \alpha)q]$. As a result, we may write

$$S_{h,t}(Q) = \sum_{\substack{(a,b) \in Q\mathcal{T} \\ \bar{b} \in I_a}} \kappa_1\left(\frac{a}{Q}, \frac{b}{Q}\right) \kappa_{h+1}\left(\frac{a}{Q}, \frac{b}{Q}\right) = \sum_{k,l=1}^{\infty} A_{k,l,t}(Q),$$

with

$$A_{k,l,t}(Q) = \#(Q(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*) \cap \{(a, b); \bar{b} \in I_a\}),$$

where $I = (0, t]$. Applying Lemma 10 in [2] to $f(a, b) = 1$, $\Omega = Q(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*)$, $A = O(Q)$, $R_1 = R_2 = O(Q/k)$, and employing (2.6) and (2.8), we infer that

$$\begin{aligned} A_{k,l,t}(Q) &= tA_{k,l}(Q) + O\left(\frac{Q}{k^2} + \frac{Q^{3/2+\varepsilon}}{k}\right) \\ &= \frac{6tQ^2}{\pi^2} \text{area}(\mathcal{T}_k^* \cap T^{-h}\mathcal{T}_l^*) + O\left(\frac{Q^{3/2+\varepsilon}}{k}\right), \end{aligned}$$

leading to

$$S_{h,t}(Q) = \frac{3tA(h)Q^2}{\pi^2} + O(Q^{3/2+\varepsilon} \log Q).$$

□

Proof of Theorem 1.2. This is similar to the proof of Theorem 1.1. We get as in (2.3)

$$\begin{aligned} T_\alpha(Q) &:= \sum_i \nu_Q(\gamma_i)^\alpha = \sum_{k=1}^{\infty} k^\alpha \#(Q\mathcal{T}_k \cap \mathbb{Z}_{\text{vis}}^2) \\ &= \sum_{k=1}^{\infty} \#(Q\mathcal{T}_k \cap \mathbb{Z}_{\text{vis}}^2) \sum_{m=1}^k (m^\alpha - (m-1)^\alpha) \\ (2.13) \quad &= \sum_{m=1}^{\infty} (m^\alpha - (m-1)^\alpha) \sum_{k=m}^{\infty} \#(Q\mathcal{T}_k \cap \mathbb{Z}_{\text{vis}}^2) \\ &= \sum_{m=1}^{2Q} (m^\alpha - (m-1)^\alpha) \#(Q\mathcal{T}_m^* \cap \mathbb{Z}_{\text{vis}}^2). \end{aligned}$$

However, Corollary 2.2 gives

$$\#(Q\mathcal{T}_m^* \cap \mathbb{Z}_{\text{vis}}^2) = \frac{6Q^2}{\pi^2} \text{area}(\mathcal{T}_m^*) + O\left(\frac{Q \log Q}{m}\right),$$

which we insert into (2.13) to get

$$T_\alpha(Q) = \frac{6Q^2}{\pi^2} \sum_{m=1}^{2Q} (m^\alpha - (m-1)^\alpha) \text{area}(\mathcal{T}_m^*) + O_\alpha(E_\alpha(Q)),$$

with $E_\alpha(Q)$ as in Theorem 1.2.

However, $Q^2 \sum_{m=Q}^{\infty} m^{\alpha-1} \text{area}(\mathcal{T}_m^*) \ll Q^2 \sum_{m=Q}^{\infty} m^{\alpha-3} \ll Q^\alpha$. Hence,

$$(2.14) \quad T_\alpha(Q) = \frac{6Q^2}{\pi^2} \sum_{m=1}^{\infty} (m^\alpha - (m-1)^\alpha) \text{area}(\mathcal{T}_m^*) + O_\alpha(E_\alpha(Q)).$$

Finally, Theorem 1.2 (i) follows from (2.14), and from

$$\begin{aligned} &\sum_{m=1}^{\infty} (m^\alpha - (m-1)^\alpha) \text{area}(\mathcal{T}_m^*) \\ &= \sum_{m=1}^{\infty} (m^\alpha - (m-1)^\alpha) \sum_{k=m}^{\infty} \text{area}(\mathcal{T}_k) \\ (2.15) \quad &= \sum_{k=1}^{\infty} \text{area}(\mathcal{T}_k) \sum_{m=1}^k (m^\alpha - (m-1)^\alpha) = \sum_{k=1}^{\infty} k^\alpha \text{area}(\mathcal{T}_k) \\ &= \iint_{\mathcal{T}} \left[\frac{1+s}{t} \right]^\alpha ds dt. \end{aligned}$$

To prove Theorem 1.2 (ii), we first write as in (2.13)

$$(2.16) \quad T_{\alpha,t}(Q) := \sum_{\gamma_i \leq t} \nu_Q(\gamma_i)^\alpha = \sum_{m=1}^{2Q} (m^\alpha - (m-1)^\alpha) \sum_{\substack{(a,b) \in Q\mathcal{T}_m^* \\ b \in I_a}} 1,$$

where $I = (0, t]$ and $I_a = [(1-t)a, a]$. Applying Lemma 10 in [2] to $f(a, b) = 1$ and $\Omega = Q\mathcal{T}_m^*$, we find

$$\sum_{\substack{(a,b) \in Q\mathcal{T}_m^* \\ b \in I_a}} 1 = \frac{6tQ^2}{\pi^2} \text{area}(\mathcal{T}_m^*) + O\left(\frac{Q^{3/2+\varepsilon}}{m}\right),$$

which gives in conjunction with (2.16) and (2.15)

$$\begin{aligned} T_{\alpha,t}(Q) &= \frac{6tQ^2}{\pi^2} \sum_{m=1}^{2Q} (m^\alpha - (m-1)^\alpha) \text{area}(\mathcal{T}_m^*) + O_\alpha(F_\alpha(Q)) \\ &= \frac{6tQ^2}{\pi^2} \iint_{\mathcal{T}} \left[\frac{1+s}{t}\right]^\alpha ds dt + O_\alpha(F_\alpha(Q)), \end{aligned}$$

where $F_\alpha(Q)$ is as in the statement of Theorem 1.2.

The equalities

$$\nu_Q(\gamma_i) = \left\lfloor \frac{Q + q_{i-1}}{q_i} \right\rfloor = k = \left\lfloor \frac{2Q + 1}{q_i} \right\rfloor - 1$$

read as

$$(q_{i-1}, q_i) \in Q\mathcal{T}_k^L \cap \mathbb{Z}_{\text{vis}}^2,$$

where

$$\mathcal{T}_k^L = \left\{ (s, t) \in \mathcal{T}_k^L; \frac{2 + 1/Q}{k + 2} < t \leq \frac{2 + 1/Q}{k + 1} \right\}.$$

Hence $L(Q, k) = \#(Q\mathcal{T}_k^L \cap \mathbb{Z}_{\text{vis}}^2)$ and by means of Corollary 2.2, we find

$$\begin{aligned} L(Q, k) &= \frac{6Q^2}{\pi^2} \text{area}(\mathcal{T}_k^L) + O\left(\frac{Q}{k} + \frac{Q \log Q}{k} + \frac{Q}{k^2}\right) \\ &= \frac{6Q^2}{\pi^2} \text{area}(\mathcal{T}_k'') + O\left(\frac{Q \log Q}{k}\right). \end{aligned}$$

The estimate on $U(Q, k)$ in (1.6) follows in a similar way.

Theorem 1.3 is derived from Lemma 10 in [2] in a similar way as above.

3. SOME NUMERICAL COMPUTATIONS

The transformation T maps each region $\mathcal{T}_k$ onto its symmetric with respect to the first bisector. That is,

$$(3.1) \quad T\mathcal{T}_k = S\mathcal{T}_k \quad \text{and} \quad T\mathcal{T}_k^* = S\mathcal{T}_k^*,$$

where S acts on $\mathcal{T}$ as

$$S(x, y) = (y, x).$$

Moreover, we notice that the inverse T^{-1} of T can be expressed as

$$(3.2) \quad T^{-1} = STS.$$

We now infer from (3.2), (3.1), and the fact that T is area-preserving that

$$(3.3) \quad \begin{aligned} \text{area}(T^h\mathcal{T}_m^* \cap \mathcal{T}_n^*) &= \text{area}(\mathcal{T}_m^* \cap T^{-h}\mathcal{T}_n^*) = \text{area}(\mathcal{T}_m^* \cap ST^hS\mathcal{T}_n^*) \\ &= \text{area}(S\mathcal{T}_m^* \cap T^hS\mathcal{T}_n^*) = \text{area}(T\mathcal{T}_m^* \cap T^hT\mathcal{T}_n^*) \\ &= \text{area}(T^h\mathcal{T}_n^* \cap \mathcal{T}_m^*). \end{aligned}$$

Thus, to evaluate $A(h)$ via (2.11), it suffices to consider $m \leq n$ only. We find the following table for the value of $\text{area}(T\mathcal{T}_m^* \cap \mathcal{T}_n^*)$

	$n = 1$	$n = 2$	$n = 3$	$n = 4$	$n \geq 5$
$m = 1$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{10}$	$\frac{2}{n(n+1)}$
$m = 2$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{30}$	$\frac{1}{210}$	0
$m = 3$	$\frac{1}{6}$	$\frac{1}{30}$	0	0	0
$m = 4$	$\frac{1}{10}$	$\frac{1}{210}$	0	0	0
$m \geq 5$	$\frac{2}{m(m+1)}$	0	0	0	0

which gives in turn

$$A(1) = 2 \cdot \frac{96}{35} = \frac{192}{35} \approx 5.4857.$$

We also find the following table for the value of $\text{area}(T^2\mathcal{T}_m^* \cap \mathcal{T}_n^*)$, which is symmetric as a result of (3.3),

	$n = 1$	$n = 2$	$n = 3$	$n = 4$	$n = 5$	$n = 6$	$n = 7$	$n = 8$	$n \geq 9$
$m = 1$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{10}$	$\frac{1}{15}$	$\frac{1}{21}$	$\frac{1}{28}$	$\frac{1}{36}$	$\frac{2}{n(n+1)}$
$m = 2$	$\frac{1}{3}$	$\frac{23}{84}$	$\frac{31}{210}$	$\frac{2}{21}$	$\frac{1}{15}$	$\frac{1}{21}$	$\frac{1}{28}$	$\frac{1}{36}$	$\frac{2}{n(n+1)}$
$m = 3$	$\frac{1}{6}$	$\frac{31}{210}$	$\frac{1}{10}$	$\frac{13}{210}$	$\frac{1}{30}$	$\frac{1}{70}$	$\frac{1}{220}$	$\frac{1}{1170}$	0
$m = 4$	$\frac{1}{10}$	$\frac{2}{21}$	$\frac{13}{210}$	$\frac{1}{42}$	$\frac{1}{231}$	0	0	0	0
$m = 5$	$\frac{1}{15}$	$\frac{1}{15}$	$\frac{1}{30}$	$\frac{1}{151}$	0	0	0	0	0
$m = 6$	$\frac{1}{21}$	$\frac{1}{21}$	$\frac{1}{70}$	0	0	0	0	0	0
$m = 7$	$\frac{1}{28}$	$\frac{1}{28}$	$\frac{1}{220}$	0	0	0	0	0	0
$m = 8$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{1170}$	0	0	0	0	0	0
$m \geq 9$	$\frac{2}{m(m+1)}$	$\frac{2}{m(m+1)}$	0	0	0	0	0	0	0

and collect

$$A(2) = 2 \cdot \frac{796727}{180180} = \frac{796727}{90090} \approx 8.8437.$$

Using the second part of Remark 2.6 and the fact that T is area-preserving, we may write for $h \geq 2$,

$$\begin{aligned} \frac{A(h)}{2} &= \sum_{m,n=1}^{c_h-1} mn \operatorname{area}(T^h \mathcal{T}_m \cap \mathcal{T}_n) + 4 \sum_{n=c_h}^{\infty} n \operatorname{area}(\mathcal{T}_n) \\ &= \sum_{m,n=1}^{c_h-1} mn \operatorname{area}(T^h \mathcal{T}_m \cap \mathcal{T}_n) + \frac{2}{(c_h+1)(c_h+2)}. \end{aligned}$$

However, each region $T^r \mathcal{T}_m$ is a finite union of triangles with rational numbers as vertex coordinates. Thus $A(h)$ is a rational number for any $h \in \mathbb{N}^*$.

REFERENCES

1. V. AUGUSTIN, F.P. BOCA, C. COBELI, A. ZAHARESCU, *The h -spacing distribution between Farey points*, Math. Proc. Camb. Phil. Soc. 131 (2001), 23–38.
2. F.P. BOCA, C. COBELI, A. ZAHARESCU, *A conjecture of R. R. Hall on Farey points*, J. Reine Angew. Mathematik 535 (2001), 207–236.
3. F.P. BOCA, C. COBELI, A. ZAHARESCU, *On the distribution of the Farey sequence with odd denominators*, Michigan Math. J. **51** (2003), 557–573.
4. R. R. HALL, P. SHIU, *The index of a Farey sequence*, Michigan Math. J. **51** (2003), 209–223.
5. R. R. HALL, G. TENENBAUM, *On consecutive Farey arcs*, Acta Arith. 44 (1984), 397–405.

FPB AND AZ: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801, USA; FBOCA@MATH.UIUC.EDU; ZAHARESCU@MATH.UIUC.EDU

INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O. Box 1-764, RO-014700 BUCHAREST, ROMANIA

RNG: INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O. Box 1-764, RO-014700 BUCHAREST, ROMANIA; RADU.GOLOGAN@IMAR.RO